

Unusual Quantum Phase in Exactly Solvable Doubly Decorated Ising-Heisenberg Models

Jozef Strečka* and Michal Jaščur

Department of Theoretical Physics and Astrophysics, Faculty of Science,
P. J. Šafárik University, Park Angelinum 9, 040 01 Košice, Slovak Republic

Received 23 August 2002, accepted 13 September 2002

Key words Ising model, eight-vertex model, non-universal criticality, bicritical points.

PACS 05.50.+q, 05.70.Jm, 75.10.Hk

Copyright line will be provided by the publisher

The quantum Heisenberg model belongs to the most intensively studied models in statistical mechanics (see [1] and references therein), since it exhibits very interesting quantum phenomena. However, due to the mathematical complexities related to the quantum nature of the model, only few exact results have been obtained in this field so far. Owing to this fact, the main aim of this work is to introduce a new statistical model, which is exactly solvable for planar lattices and simultaneously exhibits interesting quantum properties. The model to be investigated here consists of Ising- and Heisenberg-type atoms regularly distributed on planar doubly decorated lattices, as it is depicted in Fig. 1 for the case of the square lattice. In order to derive the exact results for the model under investigation, it is very convenient to express the total Hamiltonian in the form $\hat{\mathcal{H}}_d = \sum_{k=1}^{Nq/2} \hat{\mathcal{H}}_k$, where the summation is carried out over all the bonds of the original (undecorated) lattice, N denotes the total number of the Ising-type atoms and q is the coordination number of the original (undecorated) lattice. The bond Hamiltonian $\hat{\mathcal{H}}_k$ is given by (see Fig. 1)

$$\hat{\mathcal{H}}_k = -J \left[\Delta (\hat{S}_{k1}^x \hat{S}_{k2}^x + \hat{S}_{k1}^y \hat{S}_{k2}^y) + \hat{S}_{k1}^z \hat{S}_{k2}^z \right] - J_1 (\hat{S}_{k1}^z + \hat{S}_{k2}^z) (\hat{\mu}_{k1}^z + \hat{\mu}_{k2}^z) - D \left[(\hat{S}_{k1}^z)^2 + (\hat{S}_{k2}^z)^2 \right], \quad (1)$$

where $\hat{S}_{k\alpha}^\gamma$ ($\gamma = x, y, z; \alpha = 1, 2$) represent the relevant components of the standard spin-1 operators and $\hat{\mu}_{k\alpha}^z$ those of the spin-1/2 operators. The exchange interaction J_1 couples the nearest-neighboring Ising and Heisenberg atoms and similarly, J couples nearest-neighboring Heisenberg pairs. Finally, Δ describes the anisotropy in the Heisenberg interactions and D denotes the single-ion anisotropy. Taking into account




Fig. 1 Part of the doubly decorated square lattice. Black circles represent the Ising atoms with spin 1/2 and gray ones denote the decorating Heisenberg atoms with spin 1.

* E-mail: jozkos@pobox.sk, Phone: +421 55 6222121 no.231, Fax: +421 55 6222124

the standard commutation relations for the bond Hamiltonians (i.e. $[\hat{\mathcal{H}}_i, \hat{\mathcal{H}}_k] = 0, i \neq k$), one can rewrite the partition function of the decorated model in the form

$$\mathcal{Z}_d = \text{Tr} \exp(-\beta \hat{\mathcal{H}}_d) = \text{Tr}_{\{\mu\}} \prod_{k=1}^{Nq/2} \text{Tr}_{S_{k1}} \text{Tr}_{S_{k2}} \exp(-\beta \hat{\mathcal{H}}_k). \quad (2)$$

Here, the symbol $\text{Tr}_{\{\mu\}}$ means a trace over all degrees of freedom of Ising spins, $\text{Tr}_{S_{k1}} \text{Tr}_{S_{k2}}$ denotes a trace over a couple of Heisenberg spins residing on the k -th bond and $\beta = 1/(k_B T)$. To proceed further, we introduce an extended decoration-iteration transformation [2]. Namely,

$$\text{Tr}_{S_{k1}} \text{Tr}_{S_{k2}} \exp(-\beta \hat{\mathcal{H}}_k) = A \exp(\beta R \mu_{k1}^z \mu_{k2}^z), \quad (3)$$

from which, after evaluating the l.h.s., we obtain for the transformation parameters A and R the following expressions:

$$A = \sqrt{(W_0 + W_1)(W_0 + W_2)}, \quad \beta R = 2 \ln(W_0 + W_1) - 2 \ln(W_0 + W_2), \quad (4)$$

where

$$W_0 = \exp[\beta(2D - J)] + 2 \exp[\beta(D - J/2)] \cosh \left[\frac{\beta}{2} \sqrt{(2D - J)^2 + 8(J\Delta)^2} \right], \quad (5)$$

$$W_1 = 2 \exp[\beta(2D + J)] \cosh(2\beta J_1) + 4 \exp(\beta D) \cosh(\beta J \Delta) \cosh(\beta J_1), \quad (6)$$

$$W_2 = 2 \exp[\beta(2D + J)] + 4 \exp(\beta D) \cosh(\beta J \Delta). \quad (7)$$

Finally, after substituting (3) into (2) one obtains the equation $\mathcal{Z}_d = A^{Nq/2} \mathcal{Z}_0$, which relates the partition function of doubly decorated Ising-Heisenberg model ($\mathcal{Z}_d$) to that one of the original (undecorated) spin-1/2 Ising model ($\mathcal{Z}_0$), that is exactly known for many planar lattices. Although, from this simple relation we can directly calculate all relevant physical quantities over the whole range of the parameters and temperature, we restrict ourselves in this brief report only to the analysis of the ground-state properties of the system. For this purpose, we have studied the lowtemperature behavior of the sublattice magnetization $m_A = \langle \hat{\mu}_{k\alpha}^z \rangle$, $m_B = \langle \hat{S}_{k\alpha}^z \rangle$, as well as the variations of the relevant nearest-neighbor correlation functions, namely, $q_{ii}^{zz} = \langle \hat{\mu}_{k1}^z \hat{\mu}_{k1}^z \rangle$, $q_{ih}^{zz} = \langle \hat{\mu}_{k1}^z \hat{S}_{k1}^z \rangle$, $q_{hh}^{\gamma\gamma} = \langle \hat{S}_{k1}^\gamma \hat{S}_{k2}^\gamma \rangle$ ($\gamma = x, y, z$) and $\eta = \langle (\hat{S}_{k1}^z)^2 \rangle$. Before discussing the results, it is worth noticing that the ground state of our system is universal in the sense that it does not depend on the coordination number and the space dimensionality of the lattice.

At first, we have depicted in Fig. 2a the ground-state phase diagram in the $D - J_1$ space for the system without single-ion anisotropy, i.e. $D/J = 0.0$. As one can see, three different phases can appear in the system depending on the ratio between J_1 and D . The phase denoted as O_1 represents a standard ferromagnetic phase which is usually observed in the pure Ising systems. As one can expect, in this case we have $m_A = 0.5$, $m_B = 1.0$, $q_{ii}^{zz} = 0.25$, $q_{ih}^{zz} = 0.5$, $q_{hh}^{zz} = 1.0$, $\eta = 1.0$ and $q_{hh}^{xx} = q_{hh}^{yy} = 0.0$. On the other hand, the phase denoted as O_2 exhibits some unusual features, though it is also a long-range ordered ferromagnetic phase in which all Ising spins are perfectly aligned along the easy axis (thus, $m_A = 0.5$ and $q_{ii}^{zz} = 0.25$). Contrary to the perfectly ordered Ising sublattice, the Heisenberg one behaves in a different way. Indeed one finds that $m_B = 0.5$, $q_{ih}^{zz} = 0.25$, $q_{hh}^{zz} = 0.0$ and $\eta = 0.5$. Moreover, in this phase one also observes the nonvanishing short-range order in the xy plane ($q_{hh}^{xx} = q_{hh}^{yy} = 0.5$). This rather unexpected behavior is simply explainable within the valence-bond-solid picture [3], that is schematically depicted for the decorating atoms in the inset of Fig. 2a. This is, of course, the pure quantum effect which is caused by relatively strong quantum fluctuations in the system. It is also clear that the phase O_2 represents a mixture of regularly ordered Ising spins with the quantum phase on the Heisenberg sublattice and to the best of our knowledge such a phase has not been discussed in the literature before. Finally, if the anisotropy parameter Δ becomes strong enough, a disordered phase (DP) appears in the system. In addition to the trivial fact that in this phase $m_A = m_B = 0.0$ is found, one also finds $q_{ii}^{zz} = 0.25$, $q_{ih}^{zz} = q_{ih}^{zz} = 0.0$, $q_{hh}^{zz} < 0.0$ and $q_{hh}^{xx} = q_{hh}^{yy} > 0.0$. In fact, all these findings clearly indicate that the relevant disordered quantum phase is




Fig. 2 Ground-state phase diagrams of the doubly decorated Ising-Heisenberg model. O_1 means the standard ferromagnetic phase, O_2 represents the quantum ferromagnetic phase, the properties of which can be explained within the valence-bond-solid picture. The situation is schematically depicted for the decorating atoms in the inset of Fig. 2a. The large open circles represent the atomic sites with $S = 1$, the small solid circles and arrows denote the spin $1/2$ variables. The solid line shows the valence bond. Finally, DP denotes the disordered frustrated phase and the black points mark in Fig. 2b the triple points.

strongly frustrated. It is also worth noticing that the boundary between the phase O_1 and O_2 is given by $J_1 = J(\Delta - 1)$ and that separating the phase O_2 and DP is given by $J_1 = J(0.5\sqrt{1 + 8\Delta^2} - \Delta - 0.5)$.

To complete the ground-state analysis, we have shown in Fig. 2b the phase diagram of the model in the $\Delta - D$ space for $J_1 = J$. Similarly, as in the previous case we have again observed the same phases. The phase diagram, however, becomes more interesting, since in this case we have also found two triple points, namely, $T_1 = [2 - \sqrt{2}, -\sqrt{2}]$ and $T_2 = [2 + \sqrt{2}, \sqrt{2}]$. Moreover, it is apparent that the relevant phase boundaries become more complex. Here we only state that in both cases ($D = 0$ and $D \neq 0$) all phase boundaries represent the points at which the system exhibits first-order phase transitions. The detailed analysis of the ground state properties, as well as that one of the thermal behavior of the system is being left for our future work which is now in preparation.

To conclude this work, we would like to emphasize that the Ising-Heisenberg models (including the present model) represent very useful exactly solvable statistical models that can exhibit very interesting quantum phenomena. Moreover, these models can be helpful in description of some real materials. For example, the recently synthesized compound studied in Ref. [4] has the structure of the doubly decorated honeycomb lattice.

References

- [1] S. Miyashita, *Physica A* **281**, 420 (2000).
- [2] I. Syozi, in: *Phase Transition and Critical Phenomena*, Vol. 1, Eds. C. Domb and M. S. Green, Academic Press, New York 1972 (p. 269)
- [3] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, *Phys. Rev. Lett.* **59**, 799 (1987).
- [4] Z. Trávníček, Z. Smékal, A. Escuer, and J. Marek, *New. J. Chem.* **25**, 655 (2001).